

MULTIPLIER IDEALS AND FILTERED D-MODULES

NERO BUDUR

ABSTRACT. We give a Hodge-theoretic interpretation of the multiplier ideal of an effective divisor on a smooth complex variety. More precisely, we show that the associated graded coherent sheaf with respect to the jumping-number filtration can be recovered from the smallest piece of M. Saito's Hodge filtration of the $\mathcal{D}$ -module of vanishing cycles.

1. INTRODUCTION

Let D be an effective $\mathbf{Q}$ -divisor on a nonsingular complex variety X of dimension n . The multiplier ideal $\mathcal{J}(D)$ is a subsheaf of ideals of $\mathcal{O}_X$ and measures in a subtle way the singularities of D , see [La01]. The singularities of D get "worse" if $\mathcal{J}(D)$ is smaller. The main goal of this note is to give a Hodge-theoretic interpretation of multiplier ideals. That such an interpretation is possible was hinted by [Bu03] where we proved a local relation at a point $x \in X$ between $\mathcal{J}(D)$ and the mixed Hodge structure on the cohomology the Milnor fiber of an integral divisor D at x .

The natural setting for our result is the theory of mixed Hodge modules due to M. Saito ([Sa88], [Sa90]). Since we restrict our attention to the Hodge filtration only and disregard the weight filtration and the rational structure, we end up working with filtered $\mathcal{D}_X$ -modules (M, F) . Here $\mathcal{D}_X$ is the sheaf of non-commutative rings of linear algebraic differential operators (see [Bo87]). The Hodge filtration F is always assumed here to be increasing. By the Riemann-Hilbert correspondence, M corresponds to a perverse sheaf on X . For example, the trivial mixed Hodge module $\mathbf{Q}_X^H[n]$ is represented by the filtered left $\mathcal{D}_X$ -module $(\mathcal{O}_X, F)$, where $\mathrm{Gr}_p^F = 0$ for $p \neq 0$. The corresponding perverse sheaf is the shifted trivial complex $\mathbf{Q}_X[n]$.

For a non-constant regular function $f : X \rightarrow \mathbf{C}$, one has the vanishing cycles functor ψ_f which can be defined on the abelian category of mixed Hodge modules. By definition, ψ_f corresponds to ${}^p\psi_f = \psi_f[-1]$ on the category of perverse sheaves on X . For $\alpha \in (0, 1] \cap \mathbf{Q}$, let $\psi_f^\alpha \mathcal{O}_X$

Date: May 15, 2003.

correspond to the eigenspace of the semisimple part of monodromy for the eigenvalue $\exp(-2\pi i\alpha)$. M. Saito's theory provides us with a canonical filtration F on $\psi_f^\alpha \mathcal{O}_X$ (for definitions, see the introduction of [Sa88]).

Theorem. *Let X be a smooth complex variety of dimension n . Let $f : X \rightarrow \mathbf{C}$ be a non-constant regular function and $D = f^{-1}(0)$ the corresponding effective divisor. Then for $\alpha \in (0, 1]$,*

$$(1) \quad \frac{\mathcal{J}((\alpha - \epsilon) \cdot D)}{\mathcal{J}(\alpha \cdot D)} = F_1 \psi_f^\alpha \mathcal{O}_X,$$

where $0 < \epsilon \ll 1$.

Here, F_1 is the smallest piece of the Hodge filtration of the left $\mathcal{D}_X$ -module $\psi_f \mathcal{O}_X$. The values $\alpha \in (0, 1]$ for which the left-hand side of (1) is nonzero are called jumping numbers (see [Bu03], [La01], [ELSV]). The values $\alpha \in (0, 1]$ for which the right-hand side of (1) is nonzero were considered in [Sa93]. Thus the Theorem answers a question in [ELSV] regarding the relation between the two sets of values and reproves their theorem that the jumping numbers of D are roots of the Bernstein-Sato polynomial of f up to a sign.

I thank Professor L. Ein for several discussions and comments on this work.

Acknowledgement. Professor M. Saito has also proved the Theorem of this note, see [Sa03]. Moreover, he proves that the multiplier ideals $\mathcal{J}(\alpha \cdot D)$ give the V -filtration of Malgrange and Kashiwara corresponding to f on the left $\mathcal{D}_X$ -module $\mathcal{O}_X$.

2. PROOF OF THE THEOREM

We used left $\mathcal{D}$ -modules only for the introduction. We will work, as our references do, with right $\mathcal{D}_X$ -modules. The trivial right $\mathcal{D}_X$ -module is $\omega_X = \bigwedge^n \Omega_X^1$, the sheaf of regular n -forms. Locally, the action of a vector field ξ on $w \in \omega_X$ is given by $w\xi = -\text{Lie}_\xi w$, the Lie derivative. $\mathbf{Q}_X^H[n]$ is represented by the filtered right $\mathcal{D}_X$ -module (ω_X, F) , where $\text{Gr}_p^F = 0$ for $p \neq -n$. In general, the transformation from left to right $\mathcal{D}_X$ -modules is given by

$$(M, F) \mapsto (\omega_X \otimes M, F_p = \omega_X \otimes F_{p+n}),$$

$$(w \otimes u)\xi = w\xi \otimes u - w \otimes \xi u.$$

Hence (1) is equivalent to

$$(2) \quad \omega_X \otimes_{\mathcal{O}_X} \frac{\mathcal{J}((\alpha - \epsilon) \cdot D)}{\mathcal{J}(\alpha \cdot D)} = F_{1-n} \psi_f^\alpha \omega_X.$$

Let $\mu : Y \rightarrow X$ be a log resolution of (X, D) . Recall that the multiplier ideal $\mathcal{J}(\alpha \cdot D)$ is defined for all $\alpha > 0$ by

$$\mathcal{J}(\alpha \cdot D) = \mu_* \mathcal{O}_Y(K_{Y/X} - \lfloor \alpha \cdot \mu^* D \rfloor),$$

where $K_{Y/X} = K_Y - \mu^* K_X$ and $\lfloor \cdot \rfloor$ means rounding down the coefficients in a $\mathbf{Q}$ -divisor. Put

$$\mathcal{K}_\alpha(X, D) = \frac{\mathcal{J}((\alpha - \epsilon) \cdot D)}{\mathcal{J}(\alpha \cdot D)},$$

where $0 < \epsilon \ll 1$. Then $\mathcal{J}(\alpha \cdot D)$ and $\mathcal{K}_\alpha(X, D)$ are independent of the choice of μ and

$$(3) \quad \mathcal{K}_\alpha(X, D) = \mu_* (\mathcal{O}_Y(K_{Y/X}) \otimes \mathcal{K}_\alpha(Y, \mu^* D)).$$

Put $g = f \circ \mu$. Let $\mathcal{M} = (\omega_Y, F)$ be the filtered $\mathcal{D}_Y$ -module with $Gr_p^F = 0$ for $p \neq -n$. By Theorem 2.14 of [Sa90], for $\alpha \in (0, 1]$,

$$\psi_f^\alpha H^j \mu_* \mathcal{M} = H^j \mu_* \psi_g^\alpha \mathcal{M},$$

for all $j \in \mathbf{Z}$. Here $\mu_* : D^bMHM(Y) \rightarrow D^bMHM(X)$ is the direct image functor on the bounded derived categories of mixed Hodge modules (we care only about the complexes of filtered $\mathcal{D}$ -modules), and H^j is the j -th cohomology of a complex. In particular, we have an equality of filtered $\mathcal{D}_X$ -modules

$$(4) \quad \psi_f^\alpha H^0 \mu_* \mathcal{M} = H^0 \mu_* \psi_g^\alpha \mathcal{M}.$$

We will show that (2), hence the Theorem, follows by taking F_{1-n} of both sides of (4).

Lemma 2.1. $F_{1-n} \psi_f^\alpha H^0 \mu_* \mathcal{M} = F_{1-n} \psi_f^\alpha (\omega_X, F)$, for $\alpha \in (0, 1]$.

Proof. Follows from $H^0 \mu_* \mathbf{Q}_Y^H[n] = \mathbf{Q}_X^H[n]$. $\square$

Lemma 2.2. $F_{1-n} H^0 \mu_* \psi_g^\alpha \mathcal{M} = \omega_X \otimes_{\mathcal{O}_X} \mathcal{K}_\alpha(X, D)$, for $\alpha \in (0, 1]$.

Proof. Recall the definition of $\mu_*(M, F)$. Let $i_\mu : Y \rightarrow Y \times X$ be the graph of μ . Let $p : Y \times X \rightarrow X$ be the natural projection. Then

$$\mu_*(M, F) = Rp.DR_{X \times Y/X}(i_\mu)_*(M, F).$$

where $Rp.$ is the usual derived direct image for sheaves. We put from now $p_* = H^0(Rp.)$ for the usual direct image of sheaves. Recall that $DR_{X \times Y/X}(M', F)$ is defined by

$$F_p DR_{X \times Y/X}(M', F) = \left[F_{p-n} M' \otimes \bigwedge^n \Theta_Y \longrightarrow \dots \longrightarrow F_p M' \right],$$

where $F_p M'$ sits in degree zero in the last complex, and $\Theta_Y = (\Omega_Y^1)^\vee$. The definition of $(i_\mu)_*(M, F) = (M', F)$ is the same as for $\mathcal{D}_Y$ -modules,

and all we need to know about the Hodge filtration is that $F_q M' = (i_\mu)_* F_q M$ if $q = \min\{p \mid F_p M \neq 0\}$. In this case, also $q = \min\{p \mid F_p M' \neq 0\}$.

If $(M, F) = \psi_g^\alpha(\omega_Y, F)$ and $\alpha \in (0, 1]$ is such that $M \neq 0$, then $q = 1 - n$. Hence,

$$\begin{aligned} F_{1-n} H^0 \mu_* \psi_g^\alpha(M, F) &= p_*(i_\mu)_* F_{1-n} M \\ &= \mu_* F_{1-n} M, \end{aligned}$$

where the last μ_* is the usual sheaf direct image. By Lemma 2.3, $F_{1-n} M = \omega_Y \otimes \mathcal{K}_\alpha(Y, \mu^* D)$. By (3), this proves the claim. $\square$

Lemma 2.3. *The Theorem is true if D is a simple normal crossing divisor.*

Proof. By definition, for $\alpha \in (0, 1]$,

$$\psi_f^\alpha \omega_X = \mathrm{Gr}_V^\alpha(i_f)_* \omega_X,$$

where $i_f : X \rightarrow X \times \mathbf{C}$ is the graph of f , $(i_f)_*$ is the direct image for (filtered) $\mathcal{D}_X$ -modules, and V is the decreasing filtration of Malgrange and Kashiwara. The Hodge filtration on ψ_f^α is the filtration $F[1]$, where F is induced by $(i_f)_*$. Here $F[i]_p = F_{p-i}$. In particular,

$$F_{1-n} \psi_f^\alpha \omega_X = F_{-n} \mathrm{Gr}_V^\alpha(i_f)_* \omega_X.$$

By Proposition 3.5-(3.5.1) of [Sa90] applied to $(\omega_X, F[n])$, one has that

$$\begin{aligned} F_{-n} V^\alpha(i_f)_* \omega_X &= \omega_X \otimes \mathcal{O}_X(-\lfloor (\alpha - \epsilon) \cdot D \rfloor) \\ &= \omega_X \otimes \mathcal{I}((\alpha - \epsilon) \cdot D), \end{aligned}$$

where $0 < \epsilon \ll 1$. Indeed, to apply that Proposition one only has to check locally, where X has coordinates $x_1, \dots, x_n$, that $\omega(x_i \partial_i + 1) = 0$, for $\omega = dx_1 \wedge \dots \wedge dx_n$, and $\partial_i = \partial_{x_i}$. This follows from $\mathrm{Lie}_{\partial_i}(x_i \omega) = \omega$. Hence it gives $\mu = (-1, \dots, -1)$ in the above-mentioned Proposition. $\square$

REFERENCES

- [Bo87] BOREL, A.; GRIVEL, P.; KAUP, B.; HAEFLIGER, A.; MALGRANGE, B., EHLERS, F., *Algebraic D-modules*, vol. 2 of *Perspectives in Mathematics*. Academic Press Inc., Boston, MA (1987).
- [Bu03] BUDUR, N., *On Hodge spectrum and multiplier ideals*, to appear in Math. Ann. (2003).
- [ELSV] EIN, L.; LAZARSFELD, R.; SMITH, K.E.; VAROLIN, D., *Jumping coefficients of multiplier ideals*, preprint (2003).
- [La01] LAZARSFELD, R., *Positivity in Algebraic Geometry*, preliminary draft (2001).
- [Sa03] SAITO, M., *Multiplier ideals, V-filtration and spectrum*, preprint (2003).

- [Sa93] SAITO, M. *On b-function, spectrum and rational singularity*, Math. Ann. 295, 51–74 (1993).
- [Sa90] SAITO M., *Mixed Hodge modules*, Publ. Res. Inst. Math. Sci. 26, no. 2, 221–333 (1990).
- [Sa88] SAITO M., *Modules de Hodge polarisables*, Publ. Res. Inst. Math. Sci. 24, no. 6, 849–995 (1988).

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ILLINOIS AT CHICAGO
851 SOUTH MORGAN STREET (M/C 249), CHICAGO, IL 60607-7045, USA
E-mail address: nero@math.uic.edu